

**Table 13. Reported cases of HIV infection  
Dane County, Wisconsin, 1983-2008**

	Year of report											
	1983-2008(a)		2003-2007				2008			Presumed alive		
	Cases	%	Cases	Avg(b)	%	Rate(c)	Cases	%	Rate(c)	Cases	%	Rate(c)
<b>Total</b>												
Total cases(d)	1,298	100%	277	55.4	100%	12	55	100%	12	877	100%	184
<b>Disease category</b>												
AIDS	863	66%	125	25.0	45%	5	24	44%	5	469	53%	98
Non-AIDS	435	34%	152	30.4	55%	6	31	56%	7	408	47%	86
<b>Sex</b>												
Male	1,103	85%	227	45.4	82%	19	42	76%	18	720	82%	303
Female	195	15%	50	10.0	18%	4	13	24%	5	157	18%	66
Unknown	0	0%	0	0.0	0%	-	0	0%	-	0	0%	-
<b>Race/Ethnicity</b>												
White	890	69%	171	34.2	62%	8	28	51%	7	556	63%	137
African American	306	24%	76	15.2	27%	65	20	36%	87	239	27%	1,037
Hispanic	75	6%	22	4.4	8%	17	3	5%	13	60	7%	259
American Indian	6	0%	1	0.2	0%	0	0	0%	0	3	0%	156
Asian/Pacific Islander	16	1%	6	1.2	2%	4	2	4%	9	14	2%	62
Multi-racial	3	0%	1	0.2	0%	-	0	0%	-	3	0%	-
Unknown	2	0%	0	0.0	0%	-	2	4%	-	2	0%	-
<b>Age(e)</b>												
<5	10	1%	1	0.2	0%	0	3	5%	10	1	0%	3
5-14	8	1%	2	0.4	1%	0	0	0%	0	4	0%	7
15-19	18	1%	7	1.4	3%	3	2	4%	6	6	1%	18
20-24	145	11%	33	6.6	12%	15	13	24%	29	27	3%	60
25-44	934	72%	177	35.4	64%	24	25	45%	17	373	43%	259
45+	183	14%	57	11.4	21%	7	12	22%	7	466	53%	281
Unknown	0	0%	0	0.0	0%	-	0	0%	-	0	0%	-
<b>Risk exposure</b>												
Men who have sex with men (MSM)	752	58%	151	30.2	55%	-	27	49%	-	472	54%	-
MSM and IDU	95	7%	19	3.8	7%	-	0	0%	-	61	7%	-
Injecting drug use (IDU)	129	10%	18	3.6	6%	-	3	5%	-	85	10%	-
High-risk heterosexual contact	129	10%	25	5.0	9%	-	4	7%	-	106	12%	-
Other	39	3%	2	0.4	1%	-	3	5%	-	20	2%	-
Undetermined	154	12%	62	12.4	22%	-	18	33%	-	133	15%	-
<b>Year of first report</b>	<b>Cases</b>	<b>%</b>	<b>Rate(c)</b>									
Cases reported before 1999	761	59%	-									
1999	47	4%	11									
2000	47	4%	11									
2001	55	4%	13									
2002	56	4%	13									
2003	48	4%	11									
2004	64	5%	14									
2005	39	3%	8									
2006	65	5%	14									
2007	61	5%	13									
2008	55	4%	12									

a. The first cases of HIV infection in Wisconsin were reported in 1983. Thus, these represent cumulative cases through the specified date.  
b. The average annual cases in the specified period.  
c. Cases per 100,000 population. Rates not available for risk exposure groups.  
d. Demographic and risk exposure breakdown not shown if total is less than 5 cases.  
e. Age when first diagnosed with HIV infection, except for cases presumed alive where age is age on 12/31/2008.