

**Table 25. Reported cases of HIV infection
Iowa County, Wisconsin, 1983-2008**

| | Year of report | | | | | | | | | | | |
|---------------------------------|----------------|----------|----------------|--------|------|---------|-------|------|---------|----------------|------|---------|
| | 1983-2008(a) | | 2003-2007 | | | | 2008 | | | Presumed alive | | |
| | Cases | % | Cases | Avg(b) | % | Rate(c) | Cases | % | Rate(c) | Cases | % | Rate(c) |
| Total | | | | | | | | | | | | |
| Total cases(d) | 19 | 100% | 6 | 1.2 | 100% | 5 | 1 | 100% | 4 | 12 | 100% | 51 |
| Disease category | | | | | | | | | | | | |
| AIDS | 12 | 63% | 2 | 0.4 | 33% | 0 | - | - | - | 5 | 42% | 21 |
| Non-AIDS | 7 | 37% | 4 | 0.8 | 67% | 4 | - | - | - | 7 | 58% | 30 |
| Sex | | | | | | | | | | | | |
| Male | 18 | 95% | 5 | 1.0 | 83% | 8 | - | - | - | 11 | 92% | 93 |
| Female | 1 | 5% | 1 | 0.2 | 17% | 0 | - | - | - | 1 | 8% | 8 |
| Unknown | 0 | 0% | 0 | 0.0 | 0% | - | - | - | - | 0 | 0% | - |
| Race/Ethnicity | | | | | | | | | | | | |
| White | 18 | 95% | 5 | 1.0 | 83% | 4 | - | - | - | 11 | 92% | 48 |
| African American | 0 | 0% | 0 | 0.0 | 0% | 0 | - | - | - | 0 | 0% | 0 |
| Hispanic | 0 | 0% | 0 | 0.0 | 0% | 0 | - | - | - | 0 | 0% | 0 |
| American Indian | 1 | 5% | 1 | 0.2 | 17% | 0 | - | - | - | 1 | 8% | 3,030 |
| Asian/Pacific Islander | 0 | 0% | 0 | 0.0 | 0% | 0 | - | - | - | 0 | 0% | 0 |
| Multi-racial | 0 | 0% | 0 | 0.0 | 0% | - | - | - | - | 0 | 0% | - |
| Unknown | 0 | 0% | 0 | 0.0 | 0% | - | - | - | - | 0 | 0% | - |
| Age(e) | | | | | | | | | | | | |
| <5 | 0 | 0% | 0 | 0.0 | 0% | 0 | - | - | - | 0 | 0% | 0 |
| 5-14 | 0 | 0% | 0 | 0.0 | 0% | 0 | - | - | - | 0 | 0% | 0 |
| 15-19 | 2 | 11% | 1 | 0.2 | 17% | 0 | - | - | - | 1 | 8% | 63 |
| 20-24 | 1 | 5% | 1 | 0.2 | 17% | 0 | - | - | - | 0 | 0% | 0 |
| 25-44 | 14 | 74% | 3 | 0.6 | 50% | 16 | - | - | - | 6 | 50% | 98 |
| 45+ | 2 | 11% | 1 | 0.2 | 17% | 0 | - | - | - | 5 | 42% | 51 |
| Unknown | 0 | 0% | 0 | 0.0 | 0% | - | - | - | - | 0 | 0% | - |
| Risk exposure | | | | | | | | | | | | |
| Men who have sex with men (MSM) | 13 | 68% | 3 | 0.6 | 50% | - | - | - | - | 7 | 58% | - |
| MSM and IDU | 0 | 0% | 0 | 0.0 | 0% | - | - | - | - | 0 | 0% | - |
| Injecting drug use (IDU) | 1 | 5% | 0 | 0.0 | 0% | - | - | - | - | 1 | 8% | - |
| High-risk heterosexual contact | 2 | 11% | 1 | 0.2 | 17% | - | - | - | - | 1 | 8% | - |
| Other | 1 | 5% | 1 | 0.2 | 17% | - | - | - | - | 1 | 8% | - |
| Undetermined | 2 | 11% | 1 | 0.2 | 17% | - | - | - | - | 2 | 17% | - |
| Year of first report | Cases | % | Rate(c) | | | | | | | | | |
| Cases reported before 1999 | 9 | 47% | - | | | | | | | | | |
| 1999 | 0 | 0% | 0 | | | | | | | | | |
| 2000 | 1 | 5% | 4 | | | | | | | | | |
| 2001 | 1 | 5% | 4 | | | | | | | | | |
| 2002 | 1 | 5% | 4 | | | | | | | | | |
| 2003 | 0 | 0% | 0 | | | | | | | | | |
| 2004 | 0 | 0% | 0 | | | | | | | | | |
| 2005 | 1 | 5% | 4 | | | | | | | | | |
| 2006 | 2 | 11% | 8 | | | | | | | | | |
| 2007 | 3 | 16% | 13 | | | | | | | | | |
| 2008 | 1 | 5% | 4 | | | | | | | | | |

a. The first cases of HIV infection in Wisconsin were reported in 1983. Thus, these represent cumulative cases through the specified date.
b. The average annual cases in the specified period.
c. Cases per 100,000 population. Rates not available for risk exposure groups.
d. Demographic and risk exposure breakdown not shown if total is less than 5 cases.
e. Age when first diagnosed with HIV infection, except for cases presumed alive where age is age on 12/31/2008.