

**Table 1. Reported cases of sexually transmitted disease
Kenosha County
Year: 2006**

| | <u>All STDs</u> | | <u>Chlamydia</u> | | <u>Gonorrhea</u> | | <u>Syphilis</u> | | <u>Genital HSV</u> | |
|---------------------------|-----------------|---------|------------------|---------|------------------|---------|-----------------|---------|--------------------|---------|
| | Cases | Percent | Cases | Percent | Cases | Percent | Cases | Percent | Cases | Percent |
| Total | 822 | 100.0% | 639 | 100.0% | 100 | 100.0% | 6 | 100.0% | 77 | 100.0% |
| Sex | | | | | | | | | | |
| Male | 220 | 26.8% | 159 | 24.9% | 40 | 40.0% | 3 | 50.0% | 18 | 23.4% |
| Female | 601 | 73.1% | 479 | 75.0% | 60 | 60.0% | 3 | 50.0% | 59 | 76.6% |
| Unknown | 1 | 0.1% | 1 | 0.2% | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% |
| Race | | | | | | | | | | |
| White | 428 | 52.1% | 336 | 52.6% | 39 | 39.0% | 5 | 83.3% | 48 | 62.3% |
| Black | 248 | 30.2% | 186 | 29.1% | 49 | 49.0% | 0 | 0.0% | 13 | 16.9% |
| Am. Indian | 1 | 0.1% | 0 | 0.0% | 0 | 0.0% | 1 | 16.7% | 0 | 0.0% |
| Asian/Pacific Is. | 9 | 1.1% | 8 | 1.3% | 1 | 1.0% | 0 | 0.0% | 0 | 0.0% |
| Other | 26 | 3.2% | 22 | 3.4% | 2 | 2.0% | 0 | 0.0% | 2 | 2.6% |
| Unknown | 110 | 13.4% | 87 | 13.6% | 9 | 9.0% | 0 | 0.0% | 14 | 18.2% |
| Hispanic ethnicity | | | | | | | | | | |
| Hispanic | 54 | 6.6% | 51 | 8.0% | 0 | 0.0% | 0 | 0.0% | 3 | 3.9% |
| Non Hispanic | 625 | 76.0% | 480 | 75.1% | 84 | 84.0% | 6 | 100.0% | 55 | 71.4% |
| Unknown | 143 | 17.4% | 108 | 16.9% | 16 | 16.0% | 0 | 0.0% | 19 | 24.7% |
| Age group | | | | | | | | | | |
| 0-4 | 1 | 0.1% | 0 | 0.0% | 1 | 1.0% | 0 | 0.0% | 0 | 0.0% |
| 5-9 | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% |
| 10-14 | 10 | 1.2% | 7 | 1.1% | 3 | 3.0% | 0 | 0.0% | 0 | 0.0% |
| 15-19 | 255 | 31.0% | 223 | 34.9% | 19 | 19.0% | 0 | 0.0% | 13 | 16.9% |
| 20-24 | 327 | 39.8% | 271 | 42.4% | 40 | 40.0% | 0 | 0.0% | 16 | 20.8% |
| 25-29 | 123 | 15.0% | 88 | 13.8% | 17 | 17.0% | 0 | 0.0% | 18 | 23.4% |
| 30-34 | 39 | 4.7% | 27 | 4.2% | 6 | 6.0% | 1 | 16.7% | 5 | 6.5% |
| 35-39 | 30 | 3.6% | 14 | 2.2% | 6 | 6.0% | 1 | 16.7% | 9 | 11.7% |
| 40-44 | 17 | 2.1% | 5 | 0.8% | 3 | 3.0% | 2 | 33.3% | 7 | 9.1% |
| 45-49 | 10 | 1.2% | 2 | 0.3% | 3 | 3.0% | 2 | 33.3% | 3 | 3.9% |
| 50+ | 10 | 1.2% | 2 | 0.3% | 2 | 2.0% | 0 | 0.0% | 6 | 7.8% |
| Unknown | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% |
| Year of diagnosis | | | | | | | | | | |
| 1997 | 422 | - | 306 | - | 70 | - | 3 | - | 43 | - |
| 1998 | 504 | - | 327 | - | 117 | - | 1 | - | 59 | - |
| 1999 | 531 | - | 332 | - | 133 | - | 1 | - | 65 | - |
| 2000 | 546 | - | 374 | - | 126 | - | 3 | - | 43 | - |
| 2001 | 708 | - | 511 | - | 124 | - | 6 | - | 67 | - |
| 2002 | 631 | - | 451 | - | 101 | - | 3 | - | 76 | - |
| 2003 | 645 | - | 460 | - | 100 | - | 3 | - | 82 | - |
| 2004 | 789 | - | 604 | - | 116 | - | 2 | - | 67 | - |
| 2005 | 827 | - | 580 | - | 129 | - | 6 | - | 112 | - |
| 2006 | 822 | - | 639 | - | 100 | - | 6 | - | 77 | - |