

Wisconsin sexually transmitted disease surveillance report, 2007

Douglas County

| | <u>Chlamydia*</u> | | | <u>Gonorrhea*</u> | | | <u>Syphilis*</u> | | | <u>Genital HSV*</u> | | |
|---------------------------|-------------------|---------|--------|-------------------|---------|--------|------------------|---------|--------|---------------------|---------|--------|
| | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** |
| Total | 96 | 100.0% | 218 | 33 | 100.0% | 75 | 0 | - | 0 | 20 | 100.0% | 45 |
| Sex | | | | | | | | | | | | |
| Male | 21 | 21.9% | 97 | 10 | 30.3% | 46 | - | - | - | 2 | 10.0% | 9 |
| Female | 75 | 78.1% | 336 | 23 | 69.7% | 103 | - | - | - | 18 | 90.0% | 81 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Race | | | | | | | | | | | | |
| White | 83 | 86.5% | 196 | 27 | 81.8% | 64 | - | - | - | 16 | 80.0% | 38 |
| African American | 4 | 4.2% | 930 | 4 | 12.1% | 930 | - | - | - | 0 | 0.0% | 0 |
| American Indian | 2 | 2.1% | 216 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| Asian/Pacific Islander | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 1 | 5.0% | 229 |
| Other | 1 | 1.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Unknown | 6 | 6.3% | - | 2 | 6.1% | - | - | - | - | 3 | 15.0% | - |
| Hispanic ethnicity | | | | | | | | | | | | |
| Hispanic | 3 | 3.1% | 701 | 1 | 3.0% | 234 | - | - | - | 0 | 0.0% | 0 |
| Non Hispanic | 78 | 81.3% | 179 | 24 | 72.7% | 55 | - | - | - | 15 | 75.0% | 34 |
| Unknown | 15 | 15.6% | - | 8 | 24.2% | - | - | - | - | 5 | 25.0% | - |
| Age group | | | | | | | | | | | | |
| 0-4 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| 5-9 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| 10-14 | 2 | 2.1% | 71 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| 15-19 | 29 | 30.2% | 1002 | 9 | 27.3% | 311 | - | - | - | 3 | 15.0% | 104 |
| 20-24 | 41 | 42.7% | 1231 | 15 | 45.5% | 450 | - | - | - | 6 | 30.0% | 180 |
| 25-29 | 16 | 16.7% | 492 | 7 | 21.2% | 215 | - | - | - | 3 | 15.0% | 92 |
| 30-34 | 5 | 5.2% | 196 | 1 | 3.0% | 39 | - | - | - | 1 | 5.0% | 39 |
| 35-39 | 3 | 3.1% | 115 | 0 | 0.0% | 0 | - | - | - | 2 | 10.0% | 76 |
| 40-44 | 0 | 0.0% | 0 | 1 | 3.0% | 30 | - | - | - | 0 | 0.0% | 0 |
| 45-49 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 2 | 10.0% | 56 |
| 50+ | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 3 | 15.0% | 20 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |

* Demographic breakdown not available for diseases with less than or equal to 10 cases reported.

** Reported cases per 100,000 population.