

Binge drinking, high school students

Health Priority C: Alcohol and Other Substance Use and Addiction

Objective C2a: Reduce the percentage of youth who report binge drinking in the past 30 days to 26.7 percent.

2010 Target: 26.7%

Indicator: Binge drinking, high school students

Percent of Wisconsin High School Students Who Engaged in Binge Drinking in the Past 30 Days

| Year (N) | Total | | Males | | Females | |
|-------------|---------|-----|---------|-----|---------|-----|
| | Percent | +/- | Percent | +/- | Percent | +/- |
| 1999 | 34% | 3% | 39% | 4% | 29% | 4% |
| (N) | 1,336 | | 682 | | 648 | |
| 2001 | 34% | 2% | 37% | 3% | 31% | 3% |
| (N) | 2,120 | | 1,031 | | 1,084 | |
| 2003 | 28% | 2% | 30% | 3% | 26% | 3% |
| (N) | 2,121 | | 1,019 | | 1,097 | |
| 2005 | 31% | 2% | 33% | 3% | 29% | 3% |
| (N) | 2,389 | | 1,163 | | 1,219 | |

Source: Wisconsin Youth Risk Behavior Survey, Department of Public Instruction.

Note: Binge drinking is defined as consuming five or more alcoholic drinks on one occasion.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation.

Percent of Wisconsin High School Students Who Engaged in Binge Drinking in the Past 30 Days, by Race/Ethnicity

| Years (N) | Total | | Afr. Am.* | | Am. Ind.* | | Asian* | | Hispanic | | White* | |
|--------------|---------|-----|-----------|-----|-----------|-----|---------|-----|----------|-----|---------|-----|
| | Percent | +/- | Percent | +/- | Percent | +/- | Percent | +/- | Percent | +/- | Percent | +/- |
| 1999-2003 | 32% | 1% | 16% | 5% | 48% | 11% | 21% | 6% | 36% | 9% | 34% | 2% |
| (N) | 5,577 | | 397 | | 104 | | 213 | | 154 | | 4,379 | |
| 2001-2005 | 31% | 1% | 16% | 4% | 45% | 9% | 25% | 6% | 30% | 7% | 33% | 1% |
| (N) | 6,630 | | 468 | | 136 | | 272 | | 203 | | 5,219 | |

Source: Wisconsin Youth Risk Behavior Survey, Department of Public Instruction.

Note: Binge drinking is defined as consuming five or more alcoholic drinks on one occasion.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation.

* Non-Hispanic

Marijuana use, high school students

Health Priority C: Alcohol and Other Substance Use and Addiction

Objective C2b: Reduce the percentage of youth who report using marijuana in the past 30 days to 20.7 percent.

2010 Target: 20.7%

Indicator: Marijuana use, high school students

Percent of Wisconsin High School Students Who Used Marijuana in the Past 30 Days

| Year (N) | Total | | Males | | Females | |
|-------------|---------|-----|---------|-----|---------|-----|
| | Percent | +/- | Percent | +/- | Percent | +/- |
| 1999 | 22% | 2% | 23% | 3% | 20% | 3% |
| (N) | 1,336 | | 682 | | 648 | |
| 2001 | 25% | 2% | 28% | 3% | 22% | 3% |
| (N) | 2,120 | | 1,031 | | 1,084 | |
| 2003 | 22% | 2% | 24% | 3% | 19% | 3% |
| (N) | 2,121 | | 1,019 | | 1,097 | |
| 2005 | 16% | 2% | 16% | 2% | 16% | 2% |
| (N) | 2,389 | | 1,163 | | 1,219 | |

Source: Wisconsin Youth Risk Behavior Survey, Department of Public Instruction.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation.

Percent of Wisconsin High School Students Who Used Marijuana in the Past 30 Days, by Race/Ethnicity

| Years (N) | Total | | Afr. Am.* | | Am. Ind.* | | Asian* | | Hispanic | | White* | |
|--------------|---------|-----|-----------|-----|-----------|-----|---------|-----|----------|-----|---------|-----|
| | Percent | +/- | Percent | +/- | Percent | +/- | Percent | +/- | Percent | +/- | Percent | +/- |
| 1999-2003 | 23% | 1% | 30% | 6% | 41% | 11% | 14% | 5% | 27% | 8% | 22% | 1% |
| (N) | 5,577 | | 397 | | 104 | | 213 | | 154 | | 4,379 | |
| 2001-2005 | 21% | 1% | 31% | 6% | 35% | 9% | 15% | 5% | 25% | 6% | 20% | 1% |
| (N) | 6,630 | | 468 | | 136 | | 272 | | 203 | | 6,630 | |

Source: Wisconsin Youth Risk Behavior Survey, Department of Public Instruction.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation.

* Non-Hispanic

Cigarette smoking, high school students

Health Priority C: Alcohol and Other Substance Use and Addiction

Objective C2c: Reduce the percentage of youth who report using tobacco in the past 30 days to 22.4 percent.

2010 Target: 22.4%

Indicator: Cigarette smoking, high school students

Percent of Wisconsin High School Students Who Smoked Cigarettes in the Past 30 Days

| Year (N) | Total | | Males | | Females | |
|-------------|---------|-----|---------|-----|---------|-----|
| | Percent | +/- | Percent | +/- | Percent | +/- |
| 1999 | 38% | 3% | 38% | 4% | 38% | 4% |
| (N) | 1,336 | | 682 | | 648 | |
| 2001 | 33% | 2% | 29% | 3% | 37% | 3% |
| (N) | 2,120 | | 1,031 | | 1,084 | |
| 2003 | 24% | 2% | 23% | 3% | 24% | 3% |
| (N) | 2,121 | | 1,019 | | 1,097 | |
| 2005 | 23% | 2% | 24% | 3% | 22% | 2% |
| (N) | 2,389 | | 1,163 | | 1,219 | |

Source: Wisconsin Youth Risk Behavior Survey, Department of Public Instruction.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation.

Percent of Wisconsin High School Students Who Smoked Cigarettes in the Past 30 Days, by Race/Ethnicity

| Years (N) | Total | | Afr. Am.* | | Am. Ind.* | | Asian* | | Hispanic | | White* | |
|--------------|---------|-----|-----------|-----|-----------|-----|---------|-----|----------|-----|---------|-----|
| | Percent | +/- | Percent | +/- | Percent | +/- | Percent | +/- | Percent | +/- | Percent | +/- |
| 1999-2003 | 31% | 1% | 17% | 5% | 53% | 12% | 26% | 7% | 37% | 10% | 32% | 2% |
| (N) | 5,577 | | 397 | | 104 | | 213 | | 154 | | 4,379 | |
| 2001-2005 | 26% | 1% | 15% | 4% | 44% | 9% | 25% | 6% | 28% | 7% | 27% | 1% |
| (N) | 6,630 | | 468 | | 136 | | 272 | | 203 | | 5,219 | |

Source: Wisconsin Youth Risk Behavior Survey, Department of Public Instruction.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation.

* Non-Hispanic

First use of alcohol before age 13, high school students

Health Priority C: Alcohol and Other Substance Use and Addiction

Objective C2d: Reduce the percentage of youth who report first use of alcohol prior to age 13 to 24.1 percent.

2010 Target: 24.1%

Indicator: First use of alcohol before age 13, high school students

Percent of Wisconsin High School Students Who First Used Alcohol Before Age 13

| Year (N) | Total | | Males | | Females | |
|-------------|---------|-----|---------|-----|---------|-----|
| | Percent | +/- | Percent | +/- | Percent | +/- |
| 1999 | 30% | 3% | 37% | 4% | 24% | 3% |
| (N) | 1,336 | | 682 | | 648 | |
| 2001 | 28% | 2% | 30% | 3% | 27% | 3% |
| (N) | 2,120 | | 1,031 | | 1,084 | |
| 2003 | 25% | 2% | 29% | 3% | 22% | 3% |
| (N) | 2,121 | | 1,019 | | 1,097 | |
| 2005 | 24% | 2% | 28% | 3% | 19% | 2% |
| (N) | 2,389 | | 1,163 | | 1,219 | |

Source: Wisconsin Youth Risk Behavior Survey, Department of Public Instruction.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation.

Percent of Wisconsin High School Students Who First Used Alcohol Before Age 13, by Race/Ethnicity

| Years (N) | Total | | Afr. Am.* | | Am. Ind.* | | Asian* | | Hispanic | | White* | |
|--------------|---------|-----|-----------|-----|-----------|-----|---------|-----|----------|-----|---------|-----|
| | Percent | +/- | Percent | +/- | Percent | +/- | Percent | +/- | Percent | +/- | Percent | +/- |
| 1999-2003 | 28% | 1% | 28% | 6% | 47% | 11% | 25% | 7% | 29% | 9% | 28% | 1% |
| (N) | 5,577 | | 397 | | 104 | | 213 | | 154 | | 4,379 | |
| 2001-2005 | 26% | 1% | 27% | 5% | 47% | 9% | 29% | 6% | 28% | 7% | 25% | 1% |
| (N) | 6,630 | | 468 | | 136 | | 272 | | 203 | | 5,219 | |

Source: Wisconsin Youth Risk Behavior Survey, Department of Public Instruction.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation.

* Non-Hispanic

First use of marijuana before age 13, high school students

Health Priority C: Alcohol and Other Substance Use and Addiction

Objective C2e: Reduce the percentage of youth who report first use of marijuana prior to age 13 to 8.5 percent.

2010 Target: 8.5%

Indicator: First use of marijuana before age 13, high school students

Percent of Wisconsin High School Students Who First Used Marijuana Before Age 13

| Year (N) | Total | | Males | | Females | |
|-------------|---------|-----|---------|-----|---------|-----|
| | Percent | +/- | Percent | +/- | Percent | +/- |
| 1999 | 11% | 2% | 12% | 3% | 9% | 2% |
| (N) | 1,336 | | 682 | | 648 | |
| 2001 | 8% | 1% | 10% | 2% | 7% | 2% |
| (N) | 2,120 | | 1,031 | | 1,084 | |
| 2003 | 9% | 1% | 12% | 2% | 6% | 2% |
| (N) | 2,121 | | 1,019 | | 1,019 | |
| 2005 | 7% | 1% | 9% | 2% | 5% | 1% |
| (N) | 2,389 | | 1,163 | | 1,219 | |

Source: Wisconsin Youth Risk Behavior Survey, Department of Public Instruction.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation.

Percent of Wisconsin High School Students Who First Used Marijuana Before Age 13, by Race/Ethnicity

| Years (N) | Total | | Afr. Am.* | | Am. Ind.* | | Asian* | | Hispanic | | White* | |
|--------------|---------|-----|-----------|-----|-----------|-----|---------|-----|----------|-----|---------|-----|
| | Percent | +/- | Percent | +/- | Percent | +/- | Percent | +/- | Percent | +/- | Percent | +/- |
| 1999-2003 | 9% | 1% | 20% | 5% | 29% | 10% | 9% | 4% | 17% | 7% | 8% | 1% |
| (N) | 5,577 | | 397 | | 104 | | 213 | | 154 | | 4,379 | |
| 2001-2005 | 8% | 1% | 20% | 5% | 24% | 8% | 10% | 4% | 11% | 5% | 6% | 1% |
| (N) | 6,630 | | 468 | | 136 | | 272 | | 203 | | 5,219 | |

Source: Wisconsin Youth Risk Behavior Survey, Department of Public Instruction.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation.

* Non-Hispanic