

## Ever had sexual intercourse, high school students

Health Priority F: High-Risk Sexual Behavior

Objective F1a: By 2010, 30% or less of Wisconsin high school youth will report ever having had sexual intercourse.

2010 Target: Less than or equal to 30%

Indicator: Ever had sexual intercourse, high school students

### Percent of Wisconsin High School Students Who Have Ever Had Sexual Intercourse

Year (N)	Total		Males		Females	
	Percent	+/-	Percent	+/-	Percent	+/-
1999	42%	3%	41%	4%	42%	4%
(N)	1,336		682		648	
2001	39%	2%	35%	3%	44%	3%
(N)	2,120		1,031		1,084	
2003	37%	2%	36%	3%	38%	3%
(N)	2,121		1,019		1,097	
2005	40%	2%	40%	3%	40%	3%
(N)	2,389		1,163		1,219	

Source: Wisconsin Youth Risk Behavior Survey, Wisconsin Department of Public Instruction.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation.

### Percent of Wisconsin High School Students Who Have Ever Had Sexual Intercourse, by Race/Ethnicity

Years (N)	Total		Afr. Am.*		Am. Ind.*		Asian*		Hispanic		White*	
	Percent	+/-	Percent	+/-	Percent	+/-	Percent	+/-	Percent	+/-	Percent	+/-
1999-2003	39%	2%	67%	6%	59%	11%	37%	9%	46%	10%	36%	2%
(N)	5,577		397		104		213		154		4,379	
2001-2005	39%	1%	67%	6%	53%	9%	30%	6%	46%	8%	36%	1%
(N)	6,630		468		136		272		203		5,219	

Source: Wisconsin Youth Risk Behavior Survey, Wisconsin Department of Public Instruction.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation.

\* Non-Hispanic

## Syphilis incidence

Health Priority F: High-Risk Sexual Behavior

Objective F3a: By 2010, the incidence of primary and secondary syphilis in Wisconsin will be 0.2 cases per 100,000 population.

2010 Target: 0.2 cases per 100,000 population

Indicator: Syphilis incidence

### Syphilis Incidence in Wisconsin, by Sex

Year	Cases per 100,000 Population	Reported Cases			
		Total	Males	Females	Unknown
2000	2.2	120	56	63	1
2001	1.3	69	33	36	0
2002	1.8	96	68	28	0
2003	1.3	70	54	16	0
2004	1.5	84	68	16	0

Source: Bureau of Communicable Diseases and Preparedness, Division of Public Health, Department of Health and Family Services.

### Syphilis Incidence in Wisconsin, by Race

Year	Cases per 100,000 Population	Reported Cases						
		Total	African American*	American Indian*	Asian*	Hispanic	White*	Other, Unknown*
2000	2.2	120	100	3	2	6	6	3
2001	1.3	69	56	0	0	7	5	1
2002	1.8	96	44	2	2	16	28	4
2003	1.3	70	27	3	0	15	22	3
2004	1.5	84	24	1	1	9	49	0

Source: Bureau of Communicable Diseases and Preparedness, Division of Public Health, Department of Health and Family Services.

\* Non-Hispanic

### Syphilis Incidence in Wisconsin, by Age

Year	Cases per 100,000 Population	Reported Cases						
		Total	Ages 0-9	Ages 10-14	Ages 15-19	Ages 20-24	Ages 25-29	Ages 30+
2000	2.2	120	0	0	4	17	18	81
2001	1.3	69	0	0	4	9	15	41
2002	1.8	96	0	0	1	15	13	67
2003	1.3	70	0	0	6	8	7	49
2004	1.5	84	0	0	8	7	11	58

Source: Bureau of Communicable Diseases and Preparedness, Division of Public Health, Department of Health and Family Services.

## Chlamydia trachomatis incidence

Health Priority F: High-Risk Sexual Behavior

Objective F3b: By 2010, the incidence of genital Chlamydia trachomatis infection in Wisconsin will be 138 cases per 100,000 population.

2010 Target: 138 cases per 100,000 population

Indicator: Chlamydia trachomatis incidence

### Chlamydia Trachomatis Incidence in Wisconsin, by Sex

Year	Cases per 100,000 Population	Reported Cases			
		Total	Males	Females	Unknown
2000	304.2	16,317	3,996	12,314	7
2001	305.0	16,553	4,266	12,239	48
2002	312.3	17,058	4,698	12,325	35
2003	323.2	17,769	4,963	12,719	87
2004	350.2	19,293	5,308	13,970	15

Source: Bureau of Communicable Diseases and Preparedness, Division of Public Health, Department of Health and Family Services.

### Chlamydia Trachomatis Incidence in Wisconsin, by Race

Year	Cases per 100,000 Population	Reported Cases						
		Total	African American*	American Indian*	Asian*	Hispanic	White*	Other, Unknown*
2000	304.2	16,317	3,605	261	156	559	4,266	7,470
2001	305.0	16,553	4,710	266	214	714	4,493	6,156
2002	312.3	17,058	5,199	273	216	839	4,652	5,879
2003	323.2	17,769	5,294	246	251	833	5,505	5,640
2004	350.2	19,293	6,208	302	284	880	5,808	5,811

Source: Bureau of Communicable Diseases and Preparedness, Division of Public Health, Department of Health and Family Services.

\* Non-Hispanic

### Chlamydia Trachomatis Incidence in Wisconsin, by Age

Year	Cases per 100,000 Population	Reported Cases						
		Total	Ages 0-9	Ages 10-14	Ages 15-19	Ages 20-24	Ages 25-29	Ages 30+
2000	304.2	16,317	22	301	6,069	5,873	2,285	1,767
2001	305.0	16,553	15	332	6,033	6,158	2,331	1,684
2002	312.3	17,058	24	341	6,306	6,299	2,392	1,696
2003	323.2	17,769	14	325	6,483	6,709	2,551	1,687
2004	350.2	19,293	18	391	6,921	7,342	2,755	1,866

Source: Bureau of Communicable Diseases and Preparedness, Division of Public Health, Department of Health and Family Services.

## Neisseria gonorrhoeae incidence

Health Priority F: High-Risk Sexual Behavior

Objective F3c: By 2010, the incidence of Neisseria gonorrhoeae infection in Wisconsin will be 63 cases per 100,000 population.

2010 Target: 63 cases per 100,000 population

Indicator: Neisseria gonorrhoeae incidence

### Neisseria Gonorrhoeae Incidence in Wisconsin, by Sex

Year	Cases per 100,000 Population	Reported Cases			
		Total	Males	Females	Unknown
2000	130.3	6,989	2,812	4,170	7
2001	112.7	6,115	2,373	3,724	18
2002	116.5	6,362	2,618	3,732	12
2003	101.8	5,598	2,317	3,259	22
2004	92.0	5,069	2,087	2,969	13

Source: Bureau of Communicable Diseases and Preparedness, Division of Public Health, Department of Health and Family Services.

### Neisseria Gonorrhoeae Incidence in Wisconsin, by Race

Year	Cases per 100,000 Population	Reported Cases						
		Total	African American*	American Indian*	Asian*	Hispanic	White*	Other, Unknown*
2000	130.3	6,989	2,644	26	19	128	734	3,438
2001	112.7	6,115	2,783	38	31	119	716	2,428
2002	116.5	6,362	3,057	60	30	145	777	2,293
2003	101.8	5,598	2,668	37	27	117	862	1,887
2004	92.0	5,069	2,547	55	36	108	838	1,485

Source: Bureau of Communicable Diseases and Preparedness, Division of Public Health, Department of Health and Family Services.

\* Non-Hispanic

### Neisseria Gonorrhoeae Incidence in Wisconsin, by Age

Year	Cases per 100,000 Population	Reported Cases						
		Total	Ages 0-9	Ages 10-14	Ages 15-19	Ages 20-24	Ages 25-30	Ages 30+
2000	130.3	6,989	13	160	2,265	2,116	982	1,333
2001	112.7	6,115	14	141	1,982	1,858	886	1,234
2002	116.5	6,362	12	148	2,140	1,943	893	1,226
2003	101.8	5,598	8	93	1,895	1,805	789	1,008
2004	92.0	5,069	2	119	1,719	1,570	723	936

Source: Bureau of Communicable Diseases and Preparedness, Division of Public Health, Department of Health and Family Services.

## HIV infection

Health Priority F: High-Risk Sexual Behavior

Objective F3d: By 2010, the incidence of human immunodeficiency virus (HIV) infection in Wisconsin will be 2.5 cases per 100,000 population.

2010 Target: 2.5 cases per 100,000 population

Indicator: HIV infection

### Incidence of HIV Infection in Wisconsin, by Sex

Year	Cases per 100,000 Population	Reported Cases		
		Total	Males	Females
2000	7.3	389	292	97
2001	6.2	336	258	78
2002	7.1	389	293	96
2003	6.6	365	285	80
2004	7.6	417	345	72

Source: Bureau of Communicable Diseases and Preparedness, Division of Public Health, Wisconsin Department of Health and Family Services.

### Incidence of HIV Infection in Wisconsin, by Race/Ethnicity

Year	Cases per 100,000 Population	Reported Cases							
		Total	African American*	American Indian*	Asian*	Hispanic	White*	More than One Race*	Unknown*
2000	7.3	389	168	5	4	44	168	-	0
2001	6.2	336	139	2	1	47	147	-	0
2002	7.1	389	143	1	2	48	195	-	0
2003	6.6	365	131	4	3	50	172	4	1
2004	7.6	417	148	4	5	52	199	5	4

Source: Bureau of Communicable Diseases and Preparedness, Division of Public Health, Wisconsin Department of Health and Family Services.

\* Non-Hispanic ethnicity. Starting in 2003, more than one race can be designated.

### Incidence of HIV Infection in Wisconsin, by Age

Year	Cases per 100,000 Population	Reported Cases						
		Total	Ages 0-9	Ages 10-14	Ages 15-19	Ages 20-24	Ages 25-29	Ages 30+
2000	7.3	389	4	0	5	43	51	286
2001	6.2	336	7	0	12	41	44	232
2002	7.1	389	5	1	14	47	66	256
2003	6.6	365	3	1	12	41	50	258
2004	7.6	417	3	1	14	50	83	266

Source: Bureau of Communicable Diseases and Preparedness, Division of Public Health, Wisconsin Department of Health and Family Services.