## **DEPARTMENT OF HEALTH SERVICES**

Division of Care and Treatment Services F-24277 (05/2024)

STATE OF WISCONSIN 42 CFR483.420(a)(2) DHS 134.31(3)(o) DHS 94.03 & 94.09 §§ 51.61(1)(g) & (h)

# **INFORMED CONSENT FOR MEDICATION**

# Dosage and / or Side Effect information last revised on 03/13/2017

Completion of this form is voluntary. If an emergency. This consent is maintained in the clien					administered without a court	order unless in			
Name – Patient / Client (Last, First MI)			ID Num		Living Unit	Date of Birth			
, Name – Individual Preparing This Form Name		Name – Staff Cor	ame – Staff Contact		Name / Telephone Number – Institution				
MEDICATION CATEGORY	MEDICATION		RECOMMENDED DAILY TOTAL DOSAGE RANGE		ANTICIPATED DOSAGE RANGE				
Antipsychotic, Antidepressant, Bipolar, Mood stabilizing Agent		edexta xtromethorphan Irobromide/quinidine sulfate)		Dextromethorphan 20 mg/quinidine 10 mg daily x 7 days, then twice daily thereafter					
The anticipated dosage range is to be individualized, may be above or below the recommended range but no medication will be administered without your informed and written consent.  Recommended daily total dosage range of manufacturer, as stated in <i>Physician's Desk Reference</i> (PDR) or another standard reference.  This medication will be administered									
Reason for Use of Psychotropic Include DSM-5 diagnosis or the diagnos				if this is 'Off-	Label' Use)				
2. Alternative mode(s) of treatment other than OR in addition to mode: Some of these would be applicable only in an inpatient environment and/or staff changes  Positive redirection and staff interaction Individual and/or group therapy Other Alternatives:									
3. Probable consequences of NOT	rosolving the	nronosod modicat	ion aro						
Impairment of Work Activities		amily Relationships			☐ Social Functioning				
Possible increase in symptoms lead Use of seclusion or restraint Limits on access to possessions Limits on personal freedoms Limit participation in treatment and Other Consequences:		ial	☐ Interve		and leisure activities nforcement authorities or others				
<b>Note:</b> These consequences may vary depending upon whether or not the individual is in an inpatient setting. It is also possible that in unusual situations, little or no adverse consequences may occur if the medications are not administered.									
						See Page 2			

Client Initial

Date \_\_\_\_\_

4. Possible side effects, warnings, and cautions associated with this medication are listed below. This is not an all-inclusive list but is representative of items of potential clinical significance to you. For more information on this medication, you may consult further with your physician or refer to a standard text, such as the PDR. As part of monitoring some of these potential side effects, your physician may order laboratory or other tests. The treatment team will closely monitor individuals who are unable to readily communicate side effects in order to enhance care and treatment.

Continued – Possible side effects, warnings, and cautions associated with this medication.

#### **Most Common Side Effects**

The most common side effects include peripheral edema, diarrhea, vomiting, asthenia, dizziness, and cough.

#### **Less Common Side Effects**

Less common side effects include prolonged QT interval, hepatitis, immune thrombocytopenia, and systemic lupus erythematosus.

#### **Rare Side Effects**

Rare side effects include dose-dependent QTc prolongation has been reported with Nuedexta and may lead to torsades de pointes dysrhythmia, with an increased risk in patients with left ventricular dysfunction or hypertrophy or with concomitant use of QT-prolonging drugs, or strong or moderate CYP3A4 inhibitors. Arrhythmias may occur, especially with concomitant use of QT-prolonging drugs, electrolyte abnormalities such as hypokalemia or hypomagnesia, bradycardia, or family history of QT abnormality.

#### Caution

Dose-dependent QTc prolongation has been reported with Nuedexta and may lead to torsades de pointes dysrhythmia, with an increased risk in patients with left ventricular dysfunction or hypertrophy or with concomitant use of QT-prolonging drugs, or strong or moderate CYP3A4 inhibitors. Arrhythmias may occur, especially with concomitant use of QT-prolonging drugs, electrolyte abnormalities such as hypokalemia or hypomagnesia, bradycardia, or family history of QT abnormality. Severe and/or fatal cases of immune-mediated thrombocytopenia, hepatitis, including granulomatous hepatitis, and Lupus-like syndrome with polyarthritis and sometimes a positive antinuclear antibody test have been reported with quinidine.

#### Warning

Nuedexta should be avoided in patients with complete atrioventricular (AV) block without implanted pacemaker, or at high risk of complete AV block. Concomitant use with drugs containing quinidine, quinine, mefloquine, drugs that both prolong the QT interval and are metabolized by CYP2D6 (e.g., thioridazine, pimozide) and MAOIs or MAOI use within 14 days should be avoided. Use of Nuedexta with SSRIs or TCAs may increase the risk of serotonin syndrome.

See standard reference text for an all-inclusive list of side effects.

Client Initial	Date	

## By my signature below, I GIVE consent for the named medication on Page 1 and anticipated dosage range. My signature also indicates that I understand the following:

- 1. I can refuse to give consent or can withdraw my consent at any time with written notification to the institution director or designee. This will not affect my right to change my decision at a later date. If I withdraw consent after a medication is started, I realize that the medication may not be discontinued immediately. Rather, it will be tapered as rapidly as medically safe and then discontinued so as to prevent an adverse medical consequence, such as seizures, due to rapid medication withdrawal.
- Questions regarding this medication can be discussed with the Interdisciplinary Team, including the physician. The staff contact person can assist in making any necessary arrangements.
- Questions regarding any behavior support plan or behavior intervention plan, which correspond with the use of the medication, can be directed to the client's social worker, case manager, or psychologist.
- I have the right to request a review at any time of my record, pursuant to § 51.30(4)(d) or § 51.30(5)(b).
- I have a legal right to file a complaint if I feel that client rights have been inappropriately restricted. The client's social worker, case manager, or agency/facility client rights specialist may be contacted for assistance.
- My consent permits the dose to be changed within the anticipated dosage range without signing another consent.
- I understand the reasons for the use of the medication, its potential risks and benefits, other alternative treatment(s), and the probable consequences that may occur if the proposed medication is not given. I have been given adequate time to study the information and find the information to be specific, accurate, and complete.
- This medication consent is for a period effective immediately and not to exceed fifteen (15) months from the date of my signature. The

need for and continued use of this medication will be reviewed at least quarte client, will be to arrive at and maintain the client at the minimum effective dos		eam. The goal, on be	half of the			
SIGNATURES		DATE	SIGNED			
Client – If Presumed Competent to Consent/Parent of Minor/Guardian (POA-HC)	Relationship to Client  Parent Guardian (F	☐ Self OA-HC)				
Staff Present at Oral Discussion	Title					
Client / Parent of Minor / Guardian (POA-HC) Comments						
As parent/guardian (POA-HC) was not available for signature, he/she was verbally informed of the information in this consent.						
Verbal Consent						
Obtained by – PRINT – Staff Name	Date Obtained	Written Consent Received ☐ Yes ☐ No				
Obtained from – PRINT – Parent / Guardian (POA-HC) Name	Date Expires	Date Received				