

Wisconsin sexually transmitted disease surveillance report, 2012

Green County

| | Chlamydia* | | | Gonorrhea* | | | Syphilis* | | | Total* | | |
|----------------------------|------------|---------|--------|------------|---------|--------|-----------|---------|--------|--------|---------|--------|
| | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** |
| Total | | | | | | | | | | | | |
| Total | 95 | 100.0% | 263 | 9 | 100.0% | 25 | 0 | - | 0 | 104 | 100.0% | 288 |
| Sex | | | | | | | | | | | | |
| Male | 23 | 24.2% | 128 | 2 | 22.2% | 11 | - | - | - | 25 | 24.0% | 140 |
| Female | 72 | 75.8% | 396 | 7 | 77.8% | 38 | - | - | - | 79 | 76.0% | 434 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Race | | | | | | | | | | | | |
| White | 85 | 89.5% | 239 | 7 | 77.8% | 20 | - | - | - | 92 | 88.5% | 259 |
| African American | 3 | 3.2% | 943 | 2 | 22.2% | 629 | - | - | - | 5 | 4.8% | 1,572 |
| American Indian | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| Asian/Pacific Islander | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| Other | 1 | 1.1% | - | 0 | 0.0% | - | - | - | - | 1 | 1.0% | - |
| Unknown | 6 | 6.3% | - | 0 | 0.0% | - | - | - | - | 6 | 5.8% | - |
| Hispanic ethnicity | | | | | | | | | | | | |
| Hispanic | 4 | 4.2% | 603 | 1 | 11.1% | 151 | - | - | - | 5 | 4.8% | 754 |
| Not Hispanic | 86 | 90.5% | 243 | 8 | 88.9% | 23 | - | - | - | 94 | 90.4% | 265 |
| Unknown | 5 | 5.3% | - | 0 | 0.0% | - | - | - | - | 5 | 4.8% | - |
| Age | | | | | | | | | | | | |
| 0-4 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| 5-9 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| 10-14 | 3 | 3.2% | 123 | 0 | 0.0% | 0 | - | - | - | 3 | 2.9% | 123 |
| 15-19 | 27 | 28.4% | 1,112 | 4 | 44.4% | 165 | - | - | - | 31 | 29.8% | 1,276 |
| 20-24 | 43 | 45.3% | 2,287 | 2 | 22.2% | 106 | - | - | - | 45 | 43.3% | 2,394 |
| 25-29 | 14 | 14.7% | 774 | 1 | 11.1% | 55 | - | - | - | 15 | 14.4% | 830 |
| 30-34 | 6 | 6.3% | 308 | 0 | 0.0% | 0 | - | - | - | 6 | 5.8% | 308 |
| 35-39 | 0 | 0.0% | 0 | 2 | 22.2% | 82 | - | - | - | 2 | 1.9% | 82 |
| 40-44 | 2 | 2.1% | 78 | 0 | 0.0% | 0 | - | - | - | 2 | 1.9% | 78 |
| 45-49 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| 50+ | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Region of residence | | | | | | | | | | | | |
| Northeastern | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Northern | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Southeastern | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Southern | 95 | 100.0% | 263 | 9 | 100.0% | 25 | - | - | - | 104 | 100.0% | 288 |
| Western | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |

* Demographic breakdown not available for diseases with less than 5 cases reported.

** Cases per 100,000 population

