

Wisconsin sexually transmitted disease surveillance report, 2015

Brown County

| | Chlamydia* | | | Gonorrhea* | | | Syphilis* | | | Total* | | |
|----------------------------|------------|---------|--------|------------|---------|--------|-----------|---------|--------|--------|---------|--------|
| | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** |
| Total | | | | | | | | | | | | |
| Total | 1,024 | 100.0% | 418 | 130 | 100.0% | 53 | 11 | 100.0% | 4 | 1,165 | 100.0% | 475 |
| Sex | | | | | | | | | | | | |
| Male | 287 | 28.0% | 235 | 69 | 53.1% | 56 | 10 | 90.9% | 8 | 366 | 31.4% | 300 |
| Female | 737 | 72.0% | 600 | 61 | 46.9% | 50 | 1 | 9.1% | 1 | 799 | 68.6% | 650 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Race | | | | | | | | | | | | |
| White | 641 | 62.6% | 283 | 55 | 42.3% | 24 | 7 | 63.6% | 3 | 703 | 60.3% | 310 |
| African American | 220 | 21.5% | 3,910 | 51 | 39.2% | 906 | 3 | 27.3% | 53 | 274 | 23.5% | 4,869 |
| American Indian | 81 | 7.9% | 1,228 | 16 | 12.3% | 243 | 0 | 0.0% | 0 | 97 | 8.3% | 1,471 |
| Asian/Pacific Islander | 29 | 2.8% | 490 | 2 | 1.5% | 34 | 0 | 0.0% | 0 | 31 | 2.7% | 524 |
| Other | 33 | 3.2% | - | 4 | 3.1% | - | 0 | 0.0% | - | 37 | 3.2% | - |
| Unknown | 20 | 2.0% | - | 2 | 1.5% | - | 1 | 9.1% | - | 23 | 2.0% | - |
| Hispanic ethnicity | | | | | | | | | | | | |
| Hispanic | 124 | 12.1% | 824 | 7 | 5.4% | 46 | 1 | 9.1% | 7 | 132 | 11.3% | 877 |
| Not Hispanic | 890 | 86.9% | 387 | 122 | 93.8% | 53 | 10 | 90.9% | 4 | 1,022 | 87.7% | 444 |
| Unknown | 10 | 1.0% | - | 1 | 0.8% | - | 0 | 0.0% | - | 11 | 0.9% | - |
| Age | | | | | | | | | | | | |
| 0-4 | 3 | 0.3% | 18 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 3 | 0.3% | 18 |
| 5-9 | 1 | 0.1% | 6 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 1 | 0.1% | 6 |
| 10-14 | 6 | 0.6% | 37 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 6 | 0.5% | 37 |
| 15-19 | 259 | 25.3% | 1,485 | 28 | 21.5% | 161 | 1 | 9.1% | 6 | 288 | 24.7% | 1,651 |
| 20-24 | 392 | 38.3% | 2,369 | 35 | 26.9% | 212 | 2 | 18.2% | 12 | 429 | 36.8% | 2,593 |
| 25-29 | 218 | 21.3% | 1,313 | 34 | 26.2% | 205 | 4 | 36.4% | 24 | 256 | 22.0% | 1,542 |
| 30-34 | 80 | 7.8% | 478 | 11 | 8.5% | 66 | 1 | 9.1% | 6 | 92 | 7.9% | 550 |
| 35-39 | 34 | 3.3% | 202 | 14 | 10.8% | 83 | 1 | 9.1% | 6 | 49 | 4.2% | 291 |
| 40-44 | 13 | 1.3% | 69 | 2 | 1.5% | 11 | 0 | 0.0% | 0 | 15 | 1.3% | 80 |
| 45-49 | 11 | 1.1% | 55 | 3 | 2.3% | 15 | 2 | 18.2% | 10 | 16 | 1.4% | 81 |
| 50+ | 6 | 0.6% | 8 | 3 | 2.3% | 4 | 0 | 0.0% | 0 | 9 | 0.8% | 12 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Region of residence | | | | | | | | | | | | |
| Northeastern | 1,024 | 100.0% | 418 | 130 | 100.0% | 53 | 11 | 100.0% | 4 | 1,165 | 100.0% | 475 |
| Northern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Southeastern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Southern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Western | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |

* Demographic breakdown not available for diseases with less than 5 cases reported.

** Cases per 100,000 population

