

Wisconsin Sexually Transmitted Infections Surveillance Report
WAUPACA COUNTY, 2021

| | Chlamydia* | | | Gonorrhea* | | | Syphilis* | | | Total* | | |
|----------------------------|------------|---------|--------|------------|---------|--------|-----------|---------|--------|--------|---------|--------|
| | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** |
| Total cases | 127 | 100.0% | 245 | 17 | 100.0% | 33 | 3 | 100.0% | 6 | 147 | 100.0% | 283 |
| Gender | | | | | | | | | | | | |
| Male | 34 | 26.8% | 129 | 10 | 58.8% | 38 | - | - | - | 46 | 31.3% | 175 |
| Female | 93 | 73.2% | 364 | 7 | 41.2% | 27 | - | - | - | 101 | 68.7% | 396 |
| Transgender Male | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Transgender Female | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Race/ethnicity*** | | | | | | | | | | | | |
| White | 83 | 65.4% | 169 | 10 | 58.8% | 20 | - | - | - | 94 | 63.9% | 191 |
| Black | 5 | 3.9% | 1,326 | 5 | 29.4% | 1,326 | - | - | - | 10 | 6.8% | 2,653 |
| Hispanic | 10 | 7.9% | 582 | - | - | - | - | - | - | 10 | 6.8% | 582 |
| Native American | - | - | - | - | - | - | - | - | - | 0 | 0.0% | 0 |
| Asian/Pacific Islander | - | - | - | - | - | - | - | - | - | 5 | 3.4% | 1,712 |
| Other | - | - | - | - | - | - | - | - | - | 6 | 4.1% | - |
| Unknown | 21 | 16.5% | - | - | - | - | - | - | - | 22 | 15.0% | - |
| Age | | | | | | | | | | | | |
| 0-4 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| 5-9 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| 10-14 | 1 | 0.8% | 33 | 0 | 0.0% | 0 | - | - | - | 1 | 0.7% | 33 |
| 15-19 | 38 | 29.9% | 1,297 | 4 | 23.5% | 137 | - | - | - | 42 | 28.6% | 1,434 |
| 20-24 | 51 | 40.2% | 2,139 | 3 | 17.6% | 126 | - | - | - | 55 | 37.4% | 2,307 |
| 25-29 | 25 | 19.7% | 936 | 3 | 17.6% | 112 | - | - | - | 28 | 19.0% | 1,048 |
| 30-34 | 5 | 3.9% | 197 | 2 | 11.8% | 79 | - | - | - | 8 | 5.4% | 314 |
| 35-39 | 3 | 2.4% | 105 | 3 | 17.6% | 105 | - | - | - | 6 | 4.1% | 210 |
| 40-44 | 3 | 2.4% | 107 | 0 | 0.0% | 0 | - | - | - | 3 | 2.0% | 107 |
| 45-49 | 1 | 0.8% | 33 | 1 | 5.9% | 33 | - | - | - | 2 | 1.4% | 67 |
| 50+ | 0 | 0.0% | 0 | 1 | 5.9% | 4 | - | - | - | 2 | 1.4% | 8 |
| Region of residence | | | | | | | | | | | | |
| Northeastern | 768 | 100.0% | 1,480 | 220 | 100.0% | 424 | 3 | 100.0% | 6 | 991 | 100.0% | 1,910 |
| Northern | - | - | - | - | - | - | - | - | - | - | - | - |
| Southeastern | - | - | - | - | - | - | - | - | - | - | - | - |
| Southern | - | - | - | - | - | - | - | - | - | - | - | - |
| Western | - | - | - | - | - | - | - | - | - | - | - | - |

*Demographic breakdown not available for diseases with less than 5 cases reported.

**Cases per 100,000 population.

*** Race/ethnicity breakdowns are not shown if the cell in the subgroup has either a value of <5 or an estimated population size (denominator) <100.

