

RESPIRATORY VIRUS SURVEILLANCE REPORT

Week 4, Ending January 30, 2021

Wisconsin Department of Health Services | Division of Public Health Bureau of Communicable Diseases | Communicable Diseases Epidemiology Section www.dhs.wisconsin.gov/dph/bcd.htm | dhsdphbcd@dhs.wi.gov





INFLUENZA LIKE ILLNESS (ILI) ACTIVITY


STATE OF WISCONSIN



REGION V OF US (WI, MN, IL, MI, OH, IN)







ILI: HIGH LEVELS - ILI: MODERATE LEVELS - ILI: BELOW BASELINE

ILI: INSUFFICIENT DATA

AT-A-GLANCE:

Predominant Viruses of the Week: Rhino/enterovirus and SARS-CoV-2 (virus causing COVID-19)

Current Alerts:

Due to the unusually low number of influenza cases nationwide, a preliminary influenza vaccine efficacy estimate could not be determined for the 2020-21 season.

Additional data on SARS-CoV-2 (the virus causing COVID-19) trends in Wisconsin can be found at:

https://www.dhs.wisconsin.gov/covid-19/data.htm

INFLUENZA-ASSOCIATED **PEDIATRIC DEATHS REPORTED:**

	Week 4, 2021	October 1, 2020 to present
Wisconsin	0	0
Nationwide	0	1




For National US influenza surveillance statistics visit: www.cdc.gov/flu/weekly/



INFLUENZA AND PNEUMONIA-ASSOCIATED MORTALITY

Influenza and Pneumonia Deaths, Wisconsin

Influenza- and pneumonia-associated deaths by influenza season year and week, Wisconsin




Influenza- and pneumonia-associated deaths by most recent influenza season week, Wisconsin, 2020-2021 season

Influenza season week	Influenza- associated deaths (I)	Pneumonia- associated deaths (P)	Percent I+P of all deaths	Baseline I+P of all deaths	Threshold I+P of all deaths
2	0	153	12.1%	5.8%	6.6%
3	1	154	12.2%	5.9%	6.7%
4 Preliminary Data	0	98	10.4%	5.9%	6.7%
Seasonal total	1	2977	11.5%	NA	NA

Data source: DPH. Office of Health Informatics



Wisconsin positive influenza results and subtypes by PCR




Cumulative number of positive influenza PCR tests by subtype October 1, 2020 to present

	A (2009 H1N1)	Influenza A: A (H3N2)	34% A (Unknown)	B (Victoria)	Influenza B: B (Yamagata)	66% B (Unknown)	Total
Total positive (n)	0	2	11	0	0	25	38
% of total positive	0%	5%	29%	0%	0%	66%	100%

WISCONSIN LABORATORY SURVEILLANCE FOR RESPIRATORY VIRUSES BY PCR

Trends in respiratory virus activity by PCR




Week 4: Ending on January 30, 2021

		Positiv	tive Positiv		ve	Influenza A							Influenza B			
Respiratory virus	Tested	(n	1)	(%)		H3N	2	2009	Unk	nown	Vict	o ria	Yamagat	a	Unknown	
Influenza	3728	1		0%		1		0		0 0		0 0			1	
Respiratory virus Tested		sted		itive n)		ositive (%) Parain		nfluenza 1	nza 1 Parainfluenza		za 2	Parainfluenza 3		Parainfluenza 4		
Parainfluenza	Parainfluenza 323		(0	09	0%		0	0		0		0			
Respirato ry vi	Respirato ry virus Test		ed I	Positiv	e (n)	e (n) Positive (9		CoV 22	CoV 229E CoV		OC 43	C43 CoV NL63			CoV HKU1	
Coronavirus (sea	Coronavirus (seasonal) 11		2	2	0.19		1%	0		0		2			0	
Respirato ry	Respirato ry virus			Tested				Pos	itive (n)			Pos	itive	(%)		
RSV					825				0				0%			
Human metapneumovirus				327				0				0%				
Rhino-enterovirus				346				27				7.8%				
Adenovirus					11	112			0				0%			

WISCONSIN STATE SUMMARY


ILI activity trend analysis by influenza season, Wisconsin




Influenza-associated hospitalizations, Wisconsin Electronic Disease Surveillance System October 1, 2020 to present

	Total		Ir	nfluenza subty	a subtype			Required		Postpartum
Age group (years)	reported (n)	A (2009 H1N1)	A (H3N2)	A (Unknown)	В	Not reported	Admitted to ICU	mechanical ventilation	Pregnant	(≤6 weeks)
<1	0	0	0	0	0	0	0	0		
1-4	0	0	0	0	0	0	0	0		
5-17	0	0	0	0	0	0	0	0		
18-49	2	0	0	0	2	0	0	0	0	0
50-64	4	0	0	2	2	0	0	0		
65+	5	0	0	1	4	0	0	0		
Total	11	0	0	3	8	0	0	0	0	0


Reported cumulative influenza-associated hospitalizations by influenza season, Wisconsin




ILI ACTIVITY TREND ANALYSIS



Northeastern Region









ILI ACTIVITY TREND ANALYSIS


Southeastern Region



Southern Region










SEASONAL INFLUENZA VACCINATION

Cumulative percentage of Wisconsin residents who received 1 or more doses of influenza vaccine, 2020-2021 influenza season



Data source: All influenza vaccination rates presented were calculated using data from the Wisconsin Immunization Registry (numerator) and Wisconsin population estimates (denominator).

Influenza vaccine composition 2020-2021:

Trivalent (three-component) egg-based vaccines are recommended to contain:

- A/Guangdong-Maonan/SWL1536/2019 (H1N1)pdm09-like virus (updated)
- A/Hong Kong/2671/2019 (H3N2)-like virus (updated)
- B/Washington/02/2019 (B/Victoria lineage)-like virus (updated)

Quadrivalent (four-component) egg-based vaccines, which protect against a second lineage of B viruses, are recommended to contain: the three recommended viruses above, plus B/Phuket/3073/2013 -like (Yamagata lineage) virus.

Cell- or recombinant-based vaccines are recommended to contain:

- A/Hawaii/70/2019 (H1N1)pdm09-like virus (updated)
- A/Hong Kong/45/2019 (H3N2)-like virus (updated)
- B/Washington/02/2019 (B/Victoria lineage)-like virus (updated)
- B/Phuket/3073/2013-like (Yamagata lineage) virus

SEASONAL INFLUENZA VACCINATION

Percentage of Wisconsin residents who received one or more doses of influenza vaccine, by age group and influenza season



Each season includes doses administered during the same time period (August 1 through May 3).

Percentage of Wisconsin residents who received one or more doses of influenza vaccine, by race and ethnicity and region, 2020-2021 influenza season

