

RESPIRATORY VIRUS SURVEILLANCE REPORT

Week 33, Ending August 21, 2021

Wisconsin Department of Health Services | Division of Public Health
Bureau of Communicable Diseases | Communicable Diseases Epidemiology Section


www.dhs.wisconsin.gov/dph/bcd.htm | dhsdphbcd@dhs.wi.gov






INFLUENZA LIKE ILLNESS (ILI) ACTIVITY


STATE OF WISCONSIN



REGION V OF US (WI, MN, IL, MI, OH, IN)



UNITED STATES



● ILI: HIGH LEVELS ● ILI: MODERATE LEVELS ● ILI: BELOW BASELINE ● ILI: INSUFFICIENT DATA

AT-A-GLANCE:

- **Predominant Viruses of the Week:**
RSV and Rhino/enterovirus are the predominant viruses this week.

INFLUENZA-ASSOCIATED PEDIATRIC DEATHS REPORTED:


| | Week 33, 2021 | October 1, 2020 to present |
|-------------------|---------------|----------------------------|
| Wisconsin | 0 | 0 |
| Nationwide | 0 | 1 |

Current Alerts:

Additional data on SARS-CoV-2 (the virus causing COVID-19) trends in Wisconsin can be found at: <https://www.dhs.wisconsin.gov/covid-19/data.htm>




For National US influenza surveillance statistics visit: www.cdc.gov/flu/weekly/



INFLUENZA AND PNEUMONIA-ASSOCIATED MORTALITY

Influenza and Pneumonia Deaths, Wisconsin

Influenza- and pneumonia-associated deaths by influenza season year and week, Wisconsin




Influenza- and pneumonia-associated deaths by most recent influenza season week, Wisconsin, 2020-2021 season

| Influenza season week | Influenza-associated deaths (I) | Pneumonia-associated deaths (P) | Percent I+P of all deaths | Baseline I+P of all deaths | Threshold I+P of all deaths |
|------------------------|---------------------------------|---------------------------------|---------------------------|----------------------------|-----------------------------|
| 31 | 0 | 44 | 4.1% | 4.4% | 5.1% |
| 32 | 0 | 66 | 6.4% | 4.3% | 5.1% |
| 33 Preliminary Data | 0 | 61 | 6.6% | 4.3% | 5.0% |
| Seasonal total | 0 | 4607 | 8.0% | NA | NA |

Data source: [DPH, Office of Health Informatics](#)

WISCONSIN LABORATORY SURVEILLANCE FOR RESPIRATORY VIRUSES BY PCR

Wisconsin positive influenza results and subtypes by PCR




Cumulative number of positive influenza PCR tests by subtype

October 1, 2020 to present

| Cumulative number of positive influenza PCR tests by subtype, September 1, 2020 to present | | | | | | | |
|--|------------------|----------|-------------|------------------|--------------|-------------|-------|
| | Influenza A: 43% | | | Influenza B: 57% | | | Total |
| | A (2009 H1N1) | A (H3N2) | A (Unknown) | B (Victoria) | B (Yamagata) | B (Unknown) | |
| Total positive (n) | 3 | 3 | 34 | 0 | 0 | 54 | 94 |

WISCONSIN LABORATORY SURVEILLANCE FOR RESPIRATORY VIRUSES BY PCR

Trends in respiratory virus activity by PCR



Week 33: Ending on August 21, 2021

| Respiratory virus | Tested | Positive (n) | Positive (%) | Influenza A | | | Influenza B | | |
|-------------------|--------|--------------|--------------|-------------|------|---------|-------------|----------|---------|
| | | | | H3N2 | 2009 | Unknown | Victoria | Yamagata | Unknown |
| Influenza | 4454 | 1 | <0.1% | 0 | 0 | 1 | 0 | 0 | 0 |


| Respiratory virus | Tested | Positive (n) | Positive (%) | Parainfluenza 1 | Parainfluenza 2 | Parainfluenza 3 | Parainfluenza 4 |
|-------------------|--------|--------------|--------------|-----------------|-----------------|-----------------|-----------------|
| Parainfluenza | 979 | 79 | 8.1% | 0 | 10 | 67 | 2 |

| Respiratory virus | Tested | Positive (n) | Positive (%) | CoV 229E | CoV OC43 | CoV NL63 | CoV HKU1 |
|------------------------|--------|--------------|--------------|----------|----------|----------|----------|
| Coronavirus (seasonal) | 126 | 0 | 0% | 0 | 0 | 0 | 0 |

| Respiratory virus | Tested | Positive (n) | Positive (%) |
|-----------------------|--------|--------------|--------------|
| RSV | 1630 | 298 | 18.3% |
| Human metapneumovirus | 978 | 2 | 0.2% |
| Rhino-enterovirus | 900 | 104 | 11.6% |
| Adenovirus | 126 | 1 | 0.8% |

WISCONSIN STATE SUMMARY


ILI activity trend analysis by influenza season, Wisconsin



Influenza-associated hospitalizations, Wisconsin Electronic Disease Surveillance System October 1, 2020 to present


| Age group (years) | Total reported (n) | Influenza subtype | | | | | Admitted to ICU | Required mechanical ventilation | Pregnant | Postpartum (≤6 weeks) |
|-------------------|--------------------|-------------------|----------|-------------|-----------|--------------|-----------------|---------------------------------|----------|-----------------------|
| | | A (2009 H1N1) | A (H3N2) | A (Unknown) | B | Not reported | | | | |
| <1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | |
| 1-4 | 1 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | | |
| 5-17 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | | |
| 18-49 | 2 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 0 |
| 50-64 | 4 | 0 | 0 | 2 | 2 | 0 | 0 | 0 | | |
| 65+ | 12 | 0 | 0 | 2 | 10 | 0 | 1 | 0 | | |
| Total | 19 | 0 | 0 | 5 | 14 | 0 | 1 | 0 | 0 | 0 |

Reported cumulative influenza-associated hospitalizations by influenza season, Wisconsin




ILI ACTIVITY TREND ANALYSIS


Wisconsin



Northeastern Region




Northern Region




ILI ACTIVITY TREND ANALYSIS


Southeastern Region



Southern Region




Western Region



SEASONAL INFLUENZA VACCINATION

Cumulative percentage of Wisconsin residents who received 1 or more doses of influenza vaccine, 2020-2021 influenza season



Data source: All influenza vaccination rates presented were calculated using data from the Wisconsin Immunization Registry (numerator) and Wisconsin population estimates (denominator).

Influenza vaccine composition 2020-2021:

Trivalent (three-component) egg-based vaccines are recommended to contain:

- A/Guangdong-Maonan/SWL1536/2019 (H1N1)pdm09-like virus (updated)
- A/Hong Kong/2671/2019 (H3N2)-like virus (updated)
- B/Washington/02/2019 (B/Victoria lineage)-like virus (updated)


Quadrivalent (four-component) egg-based vaccines, which protect against a second lineage of B viruses, are recommended to contain: the three recommended viruses above, plus B/Phuket/3073/2013-like (Yamagata lineage) virus.

Cell- or recombinant-based vaccines are recommended to contain:

- A/Hawaii/70/2019 (H1N1)pdm09-like virus (updated)
- A/Hong Kong/45/2019 (H3N2)-like virus (updated)
- B/Washington/02/2019 (B/Victoria lineage)-like virus (updated)
- B/Phuket/3073/2013-like (Yamagata lineage) virus

SEASONAL INFLUENZA VACCINATION

Percentage of Wisconsin residents who received one or more doses of influenza



Each season includes doses administered during the same time period (August 1 through May 3).

Percentage of Wisconsin residents who received one or more doses of influenza vaccine, by race and ethnicity and region, 2020-2021 influenza season

