




Respiratory Virus Surveillance Report Week 43, Ending October 26, 2024

Wisconsin Department of Health Services | Division of Public Health |
Bureau of Communicable Diseases | Communicable Diseases Epidemiology Section |
www.dhs.wisconsin.gov/dph/bcd.html | dhsdphbcd@dhs.wi.gov

Influenza-like Illness (ILI) Activity*



^{*}Wisconsin map data comes from Wisconsin DHS, regional and country-wide data use baselines calculated by CDC.

Weekly Respiratory Virus Data, At-A-Glance

Predominant virus of the week:

Rhinovirus/Enterovirus

Key Findings:

- DHS issued a health alert regarding an increase in *Mycoplasma pneumoniae* infections: www.dhs.wisconsin.gov/disease/han.htm.
- Emergency department, hospitalization, lab testing, and wastewater data show that COVID-19 activity is decreasing but remains elevated.
- Influenza and RSV are circulating at low levels.
- For updated pertussis outbreak information go to: www.dhs.wisconsin.gov/outbreaks.

Influenza-associated pediatric deaths reported:


	Week 43, 2024	Since Sep 1, 2024
Wisconsin	0	0
Nationwide	0	0

For National US influenza surveillance statistics visit: www.cdc.gov/flu/weekly/



Respiratory Virus and Pneumonia-Associated Mortality

Percent of deaths associated with influenza, RSV, COVID-19, or pneumonia by week, Vital Records $\,$




Respiratory virus and pneumonia associated deaths by most recent 3-week period, Vital Records

Season week	Pneumonia (P)	Influenza (I)	COVID-19 (C)	RSV (R)	P, I, C or R	Percent PICR of all
41	47	1	7	0	52	5%
42	56	0	19	0	67	6%
43	41	0	9	0	45	6%




Influenza-Associated Hospitalizations

Influenza-associated hospitalizations by age group, WEDSS September 1, 2024 to present



Weekly influenza-associated hospitalizations by influenza season, WEDSS



6 8

Influenza Season	Cumulative Hospitalizations Through Week 43	Entire Season
2020-2021	3	28
2021-2022	8	1021
2022-2023	66	3610
2023-2024	26	3948
2024-2025	24	-

10 12

Influenza season week

18 20 22 24

16

14

26 28

These data are preliminary and subject to change as more information is received.




42 44 46 48 50 52 2

38 40

Wisconsin Laboratory Surveillance

Wisconsin positive influenza results and subtypes by PCR, NREVSS




Cumulative number of positive influenza PCR tests by subtype, NREVSS September 1, 2024 to present

Measure	$\begin{array}{c} {\rm Influenza} \\ {\rm A(H1N1)pdm2009} \end{array}$	Influenza A(H3N2)	Influenza A Unknown	Influenza B	Total
Total positive (n)	7	20	149	73	249
% of total positive	3%	8%	60%	29%	100%



Wisconsin Laboratory Surveillance for Respiratory Viruses

Percent postivity of respiratory viruses tested by PCR, NREVSS



05/18 06/15 07/13 08/10 09/07 10/05 11/02 11/30 12/28 01/25 02/22 03/22 04/19 05/17 06/14 07/12 08/09 Week ending

Number and percent positivity of respiratory viruses tested by PCR, NREVSS Week 43, Ending on October 26, 2024


Respiratory virus	Tested	Positive (n)	Positive (%)	H3N2	2009 H1N1	A Unknown	Influenza B
Influenza	7,273	22	0.3%	1	0	12	9
Respiratory virus	Tested	Positive (n)	Positive (%)	Parainfluenza 1	Parainfluenza 2	Parainfluenza 3	Parainfluenza 4

Respiratory virus	Tested	Positive (n)	Positive (%)
Respiratory Syncytial Virus	6,200	24	0.4%
Adenovirus	990	13	1.3%
Seasonal Coronavirus	947	7	0.7%
HMPV	989	1	0.1%
Rhinovirus/Enterovirus	1,352	346	25.6%
COVID-19	8,663	550	6.3%




Respiratory Virus Activity in the Emergency Department (ED)

Percent of ED visits with a diagnosis for a respiratory virus or acute respiratory infection (ARI), NSSP




Percent of ED visits with a diagnosis for influenza by age group, NSSP






Percent of ED visits with a diagnosis for SARS-CoV-2 by age group, NSSP




Percent of ED visits with a diagnosis for RSV by age group, NSSP






Wisconsin ILI Activity

Three-week average percent of visits for ILI by influenza season, ILINET




Average percent of visits for ILI by public health region, ILINET







Northeastern Region



Northern Region




Southeastern Region






Southern Region



Western Region





Understanding the Data

Surveillance Report Description

bui vemanee reep	D C D C D C D C D C D C D C D C D C D C
Influenza-like Illness (ILI)	Patients who present to a clinician with a fever >=100 degrees F and either a cough or sore throat.
Influenza-like Illness (ILI) Activity	Using baseline (expected value data used for comparision) in each of the public health regions in Wisconsin (https://www.dhs.wisconsin.gov/lh-depts/counties/index.htm), ILI below baseline is considered low activity, ILI between baseline and threshold levels is considered moderate activity and above threshold is considered high activity. (1)
Acute Respiratory Illness (ARI)	ARI is a broad definition designed to capture all diagnoses related to respiratory illness, including SARS-CoV-2, influenza, pneumonia, and cough
Predominant virus of the week	These data are compiled from over 40 laboratories in Wisconsin that perform rt-PCR testing, and shows the viruses that have the highest percentage of positive tests.(2)
Influenza-Associated Pediatric Mortality	Deaths among children <18 years old, with influenza as the cause of associated cause of death. This is a state and nationally reportable condition. (3)
Deaths Due to Pneumonia, SARS-CoV-2, Influenza and RSV	Proportion of deaths due to pneumonia, RSV, influenza, and SARS-CoV-2 are extracted from Vital Records managed by the Office of Health Informatics through ICD-10 codes and death certificate text searches. (4)
Respiratory Viruses by PCR	A molecular laboratory method used to detect nucleic acid (DNA/RNA) in viruses, including influenza and RSV.
Influenza-Associated Hospitalizations	Patients hospitalized for >24 hours with a laboratory-identified (by rapid antigen or rt-PCR tests) influenza.(3)
Emergency Department Data	These data are from the National Syndromic Surveillance Program or NSSP. Visit information from almost all EDs in Wisconsin are reported from hospital electronic medical records to NSSP in near-real-time. Diagnoses used included the CDC Broad Acute Respiratory DD v1, the CDC COVID-Specific DD v1, CDC Influenza DD v1, and the CDC Respiratory Syncytial Virus DD v1.(5)

Additional Resources

- The CDC Influenza Homepage (https://www.cdc.gov/flu/)
- The National Respiratory and Enteric Virus Surveillance System (NREVSS) (https://www.cdc.gov/surveillance/nrevss/index.html)

Data Sources

- 1. CDC Outpatient Influenza-like Illness Surveillance Network (ILINet)
- 2. Wisconsin Laboratory Information Network and CDC National Respiratory and Enteric Virus Surveillance System (NREVSS)
- 3. Wisconsin Electronic Disease Surveillance System (WEDSS)
- 4. Division of Public Health, Office of Health Informatics, Vital Records
- 5. National Syndromic Surveillance Program (NSSP) data from ESSENCE (Electronic Surveillance System for Early Notification of Community Based Epidemics).

